arXiv:0906.3229v1 [nucl-th] 17 Jun 2009

Particle Number Fluctuations and Correlations in Nucleus-Nucleus Collisions

V.P. Konchakovski,[1, 2] M. Hauer,[2] M.I. Gorenstein,[3, 1] and E.L. Bratkovskaya[4]

[1]*Bogolyubov Institute for Theoretical Physics, Kiev, Ukraine*

[2]*Helmholtz Research School, University of Frankfurt, Frankfurt, Germany*

[3]*Frankfurt Institute for Advanced Studies, Frankfurt, Germany*

[4]*Institute für Theoretische Physik, Frankfurt, Germany*

Abstract

Particle number fluctuations and correlations in nucleus-nucleus collisions at SPS and RHIC energies are studied within the statistical hadron-resonance gas model in different statistical ensembles and in the Hadron-String-Dynamics (HSD) transport approach. Event-by-event fluctuations of the proton to pion and kaon to proton number ratios are calculated in the HSD model for the samples of most central collision events and compared with the available experimental data. The role of the experimental acceptance and centrality selection is discussed.

PACS numbers: 24.10.Lx, 24.60.Ky, 25.75.-q

Keywords:

I. INTRODUCTION

The study of event-by-event fluctuations in high energy nucleus-nucleus collisions opens new possibilities to investigate the phase transition between hadronic and partonic matter as well as the QCD critical point (see the reviews [1]). In our recent paper [2] we have presented a systematic study of the particle number fluctuations for pions $\pi = \pi^+ + \pi^-$ and kaons $K = K^+ + K^-$ and their correlations in the statistical model (SM) in different ensembles and in the Hadron-String-Dynamics (HSD) transport model [3]. In the present paper we continue this study and consider the fluctuations of the number of protons p= $p + \bar{p}$, the π-p and K-p correlations, and fluctuations of the p/π and K/p particle number ratios. The HSD results for the fluctuations in the particle number ratios are compared with NA49 data for Pb+Pb collisions at SPS energies [4] for the proton to pion ratio as well as preliminary STAR data for Au+Au collisions at RHIC energies for the proton to pion [5] and kaon to proton [6] ratios.

The paper is organized as follows. In Section II the observables for particle number fluctuations and correlations are introduced. In Section III the SM and HSD results for the scaled variances and correlation coefficients of particle number fluctuations are presented for central nucleus-nucleus collisions at SPS and RHIC energies. In Section IV the HSD transport model results for the fluctuations of the p/π and K/p ratios are presented and compared to the available data on proton to pion ratio fluctuations. The role of the centrality selection and experimental acceptance is discussed using the HSD results. A summary closes the paper in Section V.

II. OBSERVABLES FOR PARTICLE RATIO FLUCTUATIONS

We define the covariance for particle species A and B as:

$$\Delta\,(N_A, N_B) \;\equiv\; \langle \Delta N_A \Delta N_B \rangle \;=\; \langle N_A N_B \rangle \;-\; \langle N_A \rangle \langle N_B \rangle \;, \tag{1}$$

where $\langle N_A \rangle$ is the event-by-event average of the particle multiplicity N_A and $\Delta N_A \equiv N_A - \langle N_A \rangle$. The scaled variance ω_A and correlation coefficient ρ_{AB} are defined by:

$$\omega_A \;\equiv\; \frac{\Delta\,(N_A, N_A)}{\langle N_A \rangle}\;, \qquad \rho_{AB} \;\equiv\; \frac{\langle \Delta N_A \; \Delta N_B \rangle}{\left[\langle (\Delta N_A)^2 \rangle \; \langle (\Delta N_B)^2 \rangle\right]^{1/2}}\;. \tag{2}$$

Furthermore, the fluctuations in the ratio $R_{AB} \equiv N_A/N_B$ will be characterised by:

$$\sigma^2_{AB} \;\equiv\; \frac{\langle(\Delta R_{AB})^2\rangle}{\langle R_{AB}\rangle^2} \;. \tag{3}$$

One finds to second order in $\Delta N_A/\langle N_A\rangle$ and $\Delta N_B/\langle N_B\rangle$:

$$\sigma^2_{AB} \;\cong\; \frac{\omega_A}{\langle N_A\rangle} \;+\; \frac{\omega_B}{\langle N_B\rangle} \;-\; 2\rho_{AB}\left[\frac{\omega_A\omega_B}{\langle N_A\rangle\langle N_B\rangle}\right]^{1/2} \;. \tag{4}$$

We recall that the experimental data for N_A/N_B fluctuations are usually presented in terms of the so called dynamical fluctuations [7][1]

$$\sigma^{dyn}_{AB} \;\equiv\; \mathtt{sign}\left(\sigma^2_{AB} \;-\; \sigma^2_{AB,mix}\right)\left|\sigma^2_{AB} \;-\; \sigma^2_{AB,mix}\right|^{1/2} \;, \tag{5}$$

where σ^2_{AB} is defined by Eq. (4) while $\sigma^2_{AB,mix}$ corresponds to the *mixed events* background (see Ref. [2]):

$$\sigma^2_{AB,mix} \;=\; \frac{1}{\langle N_A\rangle} \;+\; \frac{1}{\langle N_B\rangle} \;. \tag{6}$$

III. PARTICLE NUMBER FLUCTUATIONS AND CORRELATIONS

In this section we present the results of the SM and HSD for the particle number fluctuations and correlations in central nucleus-nucleus collisions. The procedure of the calculations is essentially the same as in our previous paper [2]. The SM and HSD results for scaled variances of kaons $K = K^+ + K^-$, pions $\pi = \pi^+ + \pi^-$, and protons p= $p + \bar{p}$, and the correlation coefficients $\rho_{K\pi}$, $\rho_{\pi p}$, and ρ_{Kp} (defined by Eq. (2)) are presented in Tables I and II. Figures 1 and 2 show ω_p and $\rho_{p\pi}$, ρ_{Kp}, respectively. We note the results for ω_K, ω_π and $\rho_{K\pi}$ have been presented in Ref. [2].

Let us first comment on the SM results for the particle number fluctuations and correlations in different ensembles. The global charge conservation laws suppress the particle number fluctuations in the canonical ensemble (CE) in a comparison with the grand canonical ensemble (GCE). Exact energy conservation in the micro canonical ensemble (MCE) makes this suppression even stronger. This leads to the result that the MCE scaled variances ω_A become smaller than unity. The contributions to the correlation parameter ρ_{AB} in the SM stem from

---

[1] Other dynamical measures, such as Φ [8, 9] and F [10], may be also used.

$\sqrt{s_{NN}}$	T	μ_B	n_p	GCE			CE			MCE		
[GeV]	[MeV]	[MeV]	$[fm^{-3}]$	ω_p	$\rho_{p\pi}$	ρ_{Kp}	ω_p	$\rho_{p\pi}$	ρ_{Kp}	ω_p	$\rho_{p\pi}$	ρ_{Kp}
6.27	130.7	482.4	0.048	0.995	0.207	0.009	0.447	0.035	-0.096	0.446	0.090	-0.084
7.62	138.3	424.6	0.047	0.996	0.198	0.011	0.481	0.032	-0.079	0.480	0.078	-0.070
8.77	142.9	385.4	0.046	0.997	0.190	0.012	0.506	0.032	-0.068	0.505	0.066	-0.063
12.3	151.5	300.1	0.041	0.998	0.168	0.014	0.573	0.040	-0.044	0.572	0.024	-0.060
17.3	157	228.6	0.037	0.999	0.150	0.015	0.658	0.057	-0.024	0.649	-0.027	-0.069
62.4	163.1	72.7	0.027	1.000	0.119	0.017	0.938	0.106	0.012	0.859	-0.124	-0.101
130	163.6	36.1	0.026	1.000	0.116	0.017	0.984	0.113	0.016	0.888	-0.134	-0.105
200	163.7	23.4	0.026	1.000	0.115	0.017	0.993	0.114	0.017	0.894	-0.136	-0.106

TABLE I: The chemical freeze-out parameters T and μ_B for central Pb+Pb (Au+Au) collisions along the chemical freeze-out line [11]. The hadron-resonance gas model results are presented for the number density of $p+\bar{p}$, (particle number densities are the same in all statistical ensembles in the large volume limit), scaled variance ω_p, and correlation parameter $\rho_{p\pi}$ and ρ_{Kp} in the GCE, CE, and MCE. The SM results for n_π, n_K, ω_π, ω_K and $\rho_{K\pi}$ are presented in Table I of Ref. [2]

two sources: resonance decays and global conservation laws. Resonances decaying into pairs of particles of species A and B produce a corresponding correlation and yield a positive contribution to ρ_{AB}. On the other hand, the conservation laws lead to an anticorrelation and thus to negative contributions to ρ_{AB}.

As seen from Figs. 1 and 2 the HSD results for ω_p, $\rho_{p\pi}$ and ρ_{Kp} (denoted by the solid lines) are close to the CE and MCE results for low SPS energies. One may conclude that the influence of conservation laws is more stringent at low collision energies. The same conclusion follows from the HSD results for ω_π, ω_K and $\rho_{K\pi}$ (see Figs. 1 and 2 in Ref. [2]). The HSD values for ω_A and ρ_{AB} increase, however, at high collision energies and a sizeable deviation of the HSD results from those in the MCE SM is observed with increasing energies for $\sqrt{s_{NN}} > 200$ GeV.

We point out again that important aspects of the event-by-event fluctuations in nucleus-nucleus collisions are the dependence on the centrality selection and experimental acceptance. We accordingly discuss the role of these effects using HSD results for the scaled variance ω_π of the pion number fluctuations. In Fig. 3 the scaled variance ω_π (calculated within HSD) for the

$\sqrt{s_{NN}}$	HSD full acceptance								
[GeV]	$\langle N_\pi \rangle$	$\langle N_K \rangle$	$\langle N_p \rangle$	ω_π	ω_K	ω_p	$\rho_{K\pi}$	$\rho_{p\pi}$	ρ_{Kp}
6.27	612.03	43.329	181.83	0.961	1.107	0.506	-0.091	0.048	-0.137
7.62	732.11	60.801	180.03	1.077	1.141	0.526	-0.063	0.025	-0.128
8.77	823.71	75.133	179.43	1.159	1.168	0.546	-0.033	0.016	-0.129
12.3	1072.3	116.44	180.81	1.378	1.250	0.596	0.046	-0.006	-0.126
17.3	1364.6	165.52	186.97	1.619	1.348	0.641	0.126	-0.010	-0.118
62.4	2933.9	449.29	240.53	3.006	1.891	0.863	0.412	0.074	-0.029
130	4304.2	692.59	307.31	4.538	2.378	1.020	0.557	0.177	0.067
200	5204.0	861.77	352.91	5.838	2.765	1.122	0.634	0.251	0.135

TABLE II: The HSD results for the average multiplicities $\langle N_\pi \rangle$, $\langle N_K \rangle$, $\langle N_p \rangle$ and values of ω_π, ω_K, and $\rho_{K\pi}$ for central (impact parameter $b = 0$) Pb+Pb (Au+Au) collisions at different c.m. energies $\sqrt{s_{NN}}$.

full acceptance and for the experimental acceptance are shown in nucleus-nucleus collisions for zero impact parameter $b = 0$ (see the next Section for details of the experimental acceptance). Introducing the probability q of pion experimental acceptance as the ratio of an average accepted to the total multiplicities, $q = \langle N_\pi^{acc} \rangle / \langle N_\pi^{tot} \rangle$, one finds:

$$\omega_\pi^{acc} ~=~ 1 ~-~ q ~+~ q ~\omega_\pi^{full} ~. \tag{7}$$

Equation (7) connects the scaled variance ω_π^{full} in the full 4π space with ω_π^{acc} defined for the experimental acceptance. The acceptance scaling (7) assumes (see, e.g. Ref. [12]) an absence of particle correlations in momentum space. Figure 3 demonstrates that the acceptance scaling (7) underestimates the scaled variance ω_π^{acc} at RHIC energies.

The samples of collision events selected experimentally are 3.5% of most central collision events in Pb+Pb collisions at the SPS energies and 5% in Au+Au collisions at RHIC energies. Figure 4 presents the HSD results for ω_π in these samples of the most central events and their comparison with the HSD results at zero impact parameter. One finds much larger values of ω_π in the centrality selected samples than for $b = 0$. The effect is especially strong at RHIC



FIG. 1: (Color online) The SM results in the GCE, CE, and MCE ensembles and the HSD results (impact parameter $b = 0$) are presented for the scaled variance ω_p in Pb+Pb (Au+Au) collisions at different c.m. energies $\sqrt{s_{NN}}$.

energies. This can be qualitatively understood within the model of independent sources:

$$\omega_\pi \ = \ \omega_\pi^s \ + \ n_\pi \ \omega_P \ , \qquad (8)$$

where ω_π^s is the scaled variance for pions produced by one source, ω_P is the scaled variance for the fluctuations of nucleon participants, and n_π is the average pion multiplicity per participating nucleon which increases monotonously with collision energy. Collisions with zero impact parameter correspond to $\omega_P \cong 0$. Thus, ω_π^s can be approximately taken as ω_π at $b = 0$. The HSD results correspond approximately to $\omega_P \cong 0.5$ for the 3.5% most central Pb+Pb collisions at SPS energies and $\omega_P \cong 1$ for the 5% most central Au+Au collisions at RHIC energies. Please note that we used the restrictions on impact parameter b in the HSD calculations to form the samples of most central events. The results of the model of independent sources (8) for ω_π are shown by the dashed line in Fig. 4 and are close to the actual values of the HSD simulations for ω_π.



FIG. 2: (Color online) The SM results in the GCE, CE, and MCE ensembles and the HSD results (impact parameter $b = 0$) are presented for the correlation parameters $\rho_{p\pi}$ (*upper pannel*) and ρ_{Kp} (*lower panel*) in Pb+Pb (Au+Au) collisions at different c.m. energies $\sqrt{s_{NN}}$.



FIG. 3: (Color online) The HSD results for ω_π are presented for Pb+Pb (Au+Au) collisions with zero impact parameter ($b = 0$) at different c.m. energies $\sqrt{s_{NN}}$. The upper solid line corresponds to the full 4π-acceptance and the middle one to the experimental acceptance. The lower dashed line corresponds to the acceptance scaling Eq. (7).

IV. FLUCTUATIONS OF PARTICLE NUMBER RATIOS

A comparison of the SM results for fluctuations in different ensembles with the data looks problematic at present; the same is true for most other theoretical models. This is because of difficulties in implementing the experimental acceptance and centrality selection which, however, can be taken into account in the transport approach. In order to compare the HSD calculations with the measured data, the experimental cuts are applied for the simulated set of HSD events. In Fig. 5 the HSD results of σ^{dyn} (5) for the p/π and K/p ratios are shown in comparison with the experimental data by the NA49 Collaboration at the SPS [4] and the preliminary data of the STAR Collaboration at RHIC [5]. The results of the UrQMD calculations for $\sigma^{dyn}_{p\pi}$ at the SPS energies (from Ref. [4]) are also shown by the dashed line.

For the SPS energies we use the NA49 acceptance tables from Ref. [4]. For the RHIC energies we use the following cuts: in pseudorapidity, $|\eta| < 1$, and in transverse momentum,



FIG. 4: (Color online) The HSD results for ω_π for Pb+Pb (Au+Au) collisions at different c.m. energies $\sqrt{s_{NN}}$ within the full 4π-acceptance. The lower solid line corresponds to zero impact parameter ($b = 0$) and the upper one to the experimentally selected samples of collision events. The dashed line reflects the model of independent sources (8).

$0.2 < p_T < 0.6$ GeV/c for kaons and pions and $0.4 < p_T < 1$ GeV/c for protons [5, 6, 13]. We note also, that HSD results presented in Fig. 5 correspond to the centrality selection as in the experiment: the NA49 data correspond to the 3.5% most central collisions selected via veto calorimeter, whereas in the STAR experiment the 5% most central events with the highest multiplicities in the pseudorapidity range $|\eta| < 0.5$ have been selected.

At the SPS energies the HSD simulations lead to negative values of σ^{dyn} for the proton to pion ratio. This is in agreement with the NA49 data in Pb+Pb collisions. On the other hand HSD gives large positive values of $\sigma^{dyn}_{p\pi}$ at RHIC energies which strongly overestimate the preliminary STAR data for Au+Au collisions [5]. For σ^{dyn}_{Kp} only preliminary STAR data in Au+Au collisions are available [6] which demonstrate a qualitative agreement with the HSD results (Fig. 5). The HSD results for σ^{dyn}_{Kp} show a weak energy dependence in both SPS and RHIC energy regions. A peculiar feature is, however, a strong 'jump' between the SPS and RHIC values seen in the *lower panel* of Fig. 5, in the HSD calculations which is caused by the



FIG. 5: (Color online) The solid lines present the HSD results of σ^{dyn} (5) for the $(p+\bar{p})/(\pi^+ + \pi^-)$ (*upper panel*) and $(K + K^-)/(p+\bar{p})$ (*lower panel*) ratios. The data are from Refs. [4, 5, 6].

different acceptances in the SPS and RHIC measurements.

The influence of the experimental acceptance is clearly seen at 160 A GeV where a switch from the NA49 to the STAR acceptance leads to the jump in σ^{dyn}_{Kp} by 3% - lower panel of Fig. 5. On the other hand, our calculations for Pb+Pb (3.5% central) and for Au+Au (5% central) collisions - performed within the NA49 acceptance for both cases at 160 A GeV - shows a very week sensitivity of σ^{dyn}_{Kp} on the actual choice of the collision system and centrality – cf. the coincident open circle and triangle at 160 A GeV in the lower panel of Fig. 5.

V. SUMMARY AND CONCLUSIONS

We have studied the event-by-event fluctuations of the number of protons (and anti-protons), the proton-pion and proton-kaon correlations, and the fluctuations of proton to pion and kaon to proton ratios in central Pb+Pb and Au+Au collisions from low SPS up to top RHIC energies. The analysis has been performed within the statistical hadron-resonance gas model for different statistical ensembles – the grand canonical ensemble (GCE), canonical ensemble (CE), and micro-canonical ensemble (MCE) – and in the Hadron-String-Dynamics transport approach. We have found that the HSD results at SPS energies are close to those in the CE and MCE statistical model. This indicates a dominant role of resonance decays and global conservation laws at low energy nucleus-nucleus collisions. On the other hand, substantial differences in HSD and statistical model results have been observed at RHIC energies which can be attributed to non-equilibrium dynamical effects in the HSD simulations.

The HSD results for $\sigma^{dyn}_{p\pi}$ appear to be close to the NA49 data at the SPS. The data for σ^{dyn}_{Kp} in Pb+Pb collisions at the SPS energies will be available soon and allow for further insight. A comparison of the HSD results with preliminary STAR data in Au+Au collisions at RHIC energies are not fully conclusive: σ^{dyn} from HSD calculations are approximately in agreement with data [6] for kaon to proton ratio, but overestimate the experimental results [5] for proton to pion ratio. New data on event-by-event fluctuations in Au+Au at RHIC energies will help to clarify the situation.

Acknowledgements

We like to thank M. Bleicher, W. Cassing, M. Gaździcki, W. Greiner, C. Höhne, D. Kresan, M. Mitrovski, T. Schuster, R. Stock, H. Ströbele, G. Torrieri, G. Westfall and S. Wheaton for useful discussions. This work was in part supported by the Program of Fundamental Researches of the Department of Physics and Astronomy of National Academy of Sciences, Ukraine.

---

[1] H. Heiselberg, Phys. Rep. **351**, 161 (2001); S. Jeon and V. Koch, Review for Quark-Gluon Plasma 3, eds. R. C. Hwa and X.-N. Wang, World Scientific, Singapore, 430-490 (2004) [arXiv:hep-ph/0304012]; T.K. Nayak, Int. J. Mod. Phys. E **16**, 3303 (2008); V. Koch, 0810.2520 [nucl-th]; St. Mrówczyński, Acta Phys. Polon. B **40**, 1053 (2009).

[2] M. I. Gorenstein, M. Hauer, V. P. Konchakovski, E. L. Bratkovskaya Phys. Rev. C **79** 024907 (2009).

[3] W. Ehehalt and W. Cassing, Nucl. Phys. A **602**, 449 (1996); W. Cassing and E. L. Bratkovskaya, Phys. Rep. **308**, 65 (1999).

[4] C. Alt *et al.* [NA49 Collaboration], 0808.1237 [nucl-ex].

[5] G. Westfall [STAR Collaboration], poster at **Quark Matter 2009**: p/π Fluctuations in Au-Au Collisions in STAR.

[6] J. Tian [STAR Collaboration], poster at **Quark Matter 2009**: Event-by-Event p/K Fluctuations from A+A Collisions at RHIC.

[7] S. Voloshin, V. Koch, and H. Ritter, Phys. Rev. C **60**, 024901 (1999).

[8] M. Gaździcki and St. Mrówczyński, Z. Phys. C **54**, 127 (1992).

[9] St. Mrówczyński, Phys. Lett. B **439**, 6 (1998).

[10] S. Jeon and V. Koch, Phys. Rev. Lett. **83**, 5435 (1999).

[11] J. Cleymans and K. Redlich, Phys. Rev. Lett. **81**, 5284 (1998).

[12] V.V. Begun, M. Gaździcki, M.I. Gorenstein, and O.S. Zozulya, Phys. Rev. C **70**, 034901 (2004).

[13] S. Das *et al.* [STAR Collaboration], J. Phys. G **32**, S541 (2006); B.I. Abelev *et al.*, 0901.1795 [nucl-ex].